

Research Article

DOI: 10.62046/gijams.2025.v03i02.004

Serum Concentrations of Some Electrolytes (Sodium, Potassium and Bicarbonate) in Wistar Rats Fed with Carrot (*Daucus Carota*) Extract

Onyebuchi Obia^{1*}, Regina Kalio² and Elile Peace Okpara¹

¹Department of Human Physiology, faculty of Basic Medical Sciences, College of Medical Sciences, Rivers State University, Nkpolu-Oroworukwo, Port Harcourt, Nigeria.

²Department of Human Physiology, faculty of Basic Medical Sciences, College of Health Sciences, University of Port Harcourt, Nigeria.

*Corresponding Author: Onyebuchi Obia | Received: 05.02.2025 | Accepted: 15.03.2025 | Published: 17.03.2025

Abstract: Background to the study: Plasma electrolytes regulate many physiological processes necessary for cellular survival. The present study investigated the effects of carrot (*Daucus carota*) extract on serum concentrations of some electrolytes (sodium, potassium and bicarbonates) in wistar rats. **Methodology:** The study involved 28 male wistar rats separated into 4 groups of 7 rats each. Group 1 served as control and was given distilled water, whilst groups 2, 3 and 4 served as test groups and were given aqueous extract of *Daucus carota* at daily doses of 200mg/kg, 400mg/kg and 600mg/kg respectively. The experiment lasted for 28 days and thereafter the animals were sacrificed under anesthesia and blood samples collected for determination of serum potassium, sodium and bicarbonate concentrations using standard laboratory techniques. **Results and Discussion:** The results of our study showed that daily oral administration of 200mg/kg and 400mg/kg carrot extract caused a dose-dependent increase in the serum concentrations of potassium, sodium and bicarbonates, although not significantly. However, administration of 600mg/kg of carrot extract caused a significant increase in sodium concentration. Therefore, increasing the daily dosage of carrot might distort the physiological homeostatic processes with resultant significant increase in plasma sodium concentration. Excessive consumption of dietary sodium is discouraged because of risk of developing cardiovascular and other disorders. **Conclusion:** Conclusively, moderate consumption of carrot did not have any significant effect on the serum concentrations of potassium, sodium and bicarbonates and by implication did not alter the physiologic homeostatic processes. Higher doses of carrot extract significantly increased serum sodium necessitating cautious consumption of large amounts of carrot.

Keywords: Serum electrolytes, Sodium, Potassium, Bicarbonate, *Daucus carota*, Wistar rats.

Citation: Onyebuchi Obia *et al.* Serum Concentrations of Some Electrolytes (Sodium, Potassium and Bicarbonate) in Wistar Rats Fed with Carrot (*Daucus Carota*) Extract. Grn Int J Apl Med Sci, 2025 Mar-Apr 3(2): 53-56.

INTRODUCTION

Electrolytes in the blood essentially regulate many body functions; water balance, nerve and muscle function, acid-base balance, blood pressure control, maintenance of heart rate and rhythm, bone and teeth function [1]. These roles are necessary for the optimal functioning of vital organs of the body. In physiological condition, the concentrations of the different electrolytes in the blood must be maintained within their respective normal ranges. Therefore, adverse deviations from the physiologic ranges could be detrimental to cellular survival. Many of the electrolytes are derived from our diets especially fruits and other plant products [2-7]. In deficiency disorders or in conditions requiring higher demand for specific electrolytes, preformed supplements are often administered. Carrots are mainly orange coloured root vegetables [8,9] which are

consumed as food and also used in herbal preparations. Some of the health benefits of carrots have been documented in previous studies [10-13]. Although some researchers have reported significant amounts of electrolytes in a various fruits and vegetables, our study investigated the effect of carrot consumption on the serum concentrations of potassium, sodium and bicarbonates.

MATERIALS AND METHODS

The experiment was carried out at animal house of the department of Human Physiology, faculty of Basic Medical Sciences, University of Port Harcourt in the year, 2019. Ethical approval was obtained from the university of Port Harcourt Research Ethics Committee with approval number: UPH/CEREMAD/REC/MM67/012. A total of twenty

eight male wistar rats weighing 120 to 150g were purchased and acclimatized for a period of two weeks. These animals were separated into 4 groups of 7 rats each, housed in plastic cages and allowed to feed and drink *ad libitum* with Top feed Finisher mash and clean water. Their immediate environment (beddings) was changed daily, the temperature of the environment kept at normal conditions while the external environment was cleaned and disinfected regularly.

Preparation and administration of carrot extract

Mature carrot tubers were bought from Oil Mill market in Obio Akpor Local Government Area of Rivers State, Nigeria. The plant was identified at the department of Plant Science and Biotechnology, University of Port Harcourt; *Daucus carota L*, in the family; Apiceace with assigned herbarium number; UPH/C/132. The tubers were washed with water to remove soil particles. About 2kg of the fresh carrot was cut into tiny pieces and air dried for seven days. The dried carrots were blended using a blender and carefully poured into a maceration jar containing four liters of water. The mixture was allowed to macerate for 24hours after which a Whatman filter (20-25µm, pore size) was used

to get a clear filtrate. The filtrate was now poured into an evaporating dish and dried in a water bath at 45°C to obtain a semi-solid aqueous extract of *Daucus carota*. The dosages administered in the study were based on the lethal dose (LD50) of 5000mg/kg which was previously determined [14]. Following acclimatization, the wistar rats were weighed and separated into four groups of seven rats each. Group 1 served as control and was given distilled water, whilst groups 2, 3 and 4 served as test groups and were given aqueous extract of *Daucus carota* at daily doses of 200mg/kg, 400mg/kg and 600mg/kg respectively. The experiment lasted for 28 days. Thereafter the animals were sacrificed under anesthesia and blood samples collected for determination of serum potassium, sodium and bicarbonate concentrations using standard laboratory techniques.

Statistical analysis was done using SPSS software version 22.0. Results were presented in tables. Continuous variables were expressed as mean ± Standard error of mean (SEM). Statistical difference was determined using analysis of variance (ANOVA) and at $p < 0.05$.

RESULTS AND DISCUSSION

Table-1: Effect of Carrot extract on serum potassium, sodium and bicarbonate concentrations of wistar rats

Group	Potassium (mmol/l)	Sodium (mmol/l)	Bicarbonate (mmol/l)
Control	5.73±0.73	128.50±2.23	25.67±0.95
200mg/kg	6.45±0.32	129.17±3.25	27.17±0.40
400mg/kg	7.43±0.45	142.83±7.93	29.00±0.86
600mg/kg	8.70±0.40	163.00±5.72*	30.67±0.67

* Significantly different compared to control.

The results of our study showed that oral administration of carrot extract caused a dose-dependent increase in the serum concentrations of potassium and bicarbonates, although not significantly. Also the 200mg/kg and 400mg/kg of carrot extract caused a non-significant increase in sodium concentration. However, administration of 600mg/kg of carrot extract caused a significant increase in serum sodium concentration.

Despite the numerous benefits of electrolytes in the blood, the concentration and type of electrolyte consumed is of utmost importance. Most diets are enriched with sodium either to taste or as a preservative [15-17] to the extent that our daily dietary sodium requirement is often exceeded. Deficiency of plasma sodium resulting from inadequate dietary intake rarely occurs since it is richly available in our daily meals. However, hyponatremia may occur in malnutrition [18] or during conditions resulting to excessive loss of sodium [19]. Moderate consumption of carrot as shown in the present study will add just enough sodium in the plasma that is within the physiologic limit. Increasing the daily dosage of carrot up to 600mg/kg/day distorted

the physiological homeostatic processes with resultant significant increase in plasma sodium concentration. Excessive consumption of dietary sodium is discouraged because of risk of developing cardiovascular and other disorders. Therefore, carrot should be consumed moderately to obtain its benefits. Many studies support the moderate consumption of natural products for optimal benefit [20-22].

Although, some refined foods are fortified with potassium, dietary supplementation of this electrolyte is not a common practice in most food cultures. For this reason, potassium-rich foods are to be intentionally consumed to meet the daily potassium requirement. Potassium deficiency has been implicated in many diseases including cardiovascular diseases, osteoporosis and kidney stone [23-24]. Studies have shown that sodium bicarbonate composition of certain fruits and vegetables are capable of reducing dietary acid by half [25] which is a protective effect against kidney injury. The bicarbonate content of carrot can be beneficial in patients with excessive gastric acidity as it can have possible antacid potential.



CONCLUSION

Conclusively, moderate consumption of carrot did not have any significant effect on the serum concentrations of potassium, sodium and bicarbonates and by implication did not alter the physiologic homeostatic processes. Higher doses of carrot extract significantly increased serum sodium necessitating cautious consumption of large amounts of carrot.

REFERENCES

- Jomova, K., Makova, M., Alomar, S. Y., Alwasel, S. H., Nepovimova, E., Kuca, K., Rhodes, C. J., & Valko, M. (2022). Essential metals in health and disease. *Chemico-biological interactions*, 367, 110173. <https://doi.org/10.1016/j.cbi.2022.110173>
- Fernandez, M. A., & Marette, A. (2017). Potential Health Benefits of Combining Yogurt and Fruits Based on Their Probiotic and Prebiotic Properties. *Advances in nutrition (Bethesda, Md.)*, 8(1), 155S–164S. <https://doi.org/10.3945/an.115.011114>
- Qian Q. (2018). Dietary Influence on Body Fluid Acid-Base and Volume Balance: The Deleterious "Norm" Furthers and Cloaks Subclinical Pathophysiology. *Nutrients*, 10(6), 778. <https://doi.org/10.3390/nu10060778>
- Kalaycioğlu, Z., & Erim, F. B. (2019). Nitrate and Nitrites in Foods: Worldwide Regional Distribution in View of Their Risks and Benefits. *Journal of agricultural and food chemistry*, 67(26), 7205–7222. <https://doi.org/10.1021/acs.jafc.9b01194>
- Alwis, U. S., Haddad, R., Monaghan, T. F., Abrams, P., Dmochowski, R., Bower, W., Wein, A. J., Roggeman, S., Weiss, J. P., Mourad, S., Delanghe, J., & Everaert, K. (2020). Impact of food and drinks on urine production: A systematic review. *International journal of clinical practice*, 74(9), e13539. <https://doi.org/10.1111/ijcp.13539>
- Sultana, R., Alashi, A. M., Islam, K., Saifullah, M., Haque, C. E., & Aluko, R. E. (2021). Chemical composition and in vitro antioxidant properties of water-soluble extracts obtained from Bangladesh vegetables. *Journal of food biochemistry*, 45(3), e13357.
- Muhammad, A. I., Ibrahim, R. G., Adamu, S. B., Ubali, S. ., Mustapha, K., & Ibrahim, F. G. (2024). Effect of Dietary Turmeric-Ginger Combination on Serum Electrolytes Of Broiler Chicken. *Nigerian Journal of Animal Production*, 505–508.
- Carlos, J., Dias, S. (2014). Nutritional and Health Benefits of Carrots and Their Seed Extracts. *Food and Nutrition Sciences*, 5: 2147-2156.
- Bahrami, R., Ghobadi, A., Behnoud, N., & Akhtari, E. (2018). Medicinal properties of *Daucus carota* in traditional Persian medicine and modern phytotherapy. *Journal of Biochemical Technology. Special issue* (2), 107-114.
- Tanaka, T., Shnimizu, M., & Moriwaki, H. (2012). Cancer chemoprevention by carotenoids. *Molecules (Basel, Switzerland)*, 17(3), 3202–3242. <https://doi.org/10.3390/molecules17033202>
- Fiedor, J., & Burda, K. (2014). Potential role of carotenoids as antioxidants in human health and disease. *Nutrients*, 6(2), 466–488. <https://doi.org/10.3390/nu6020466>
- Bystrická, J., Kavalcová, P., Musilová, J., Vollmannová, A., Tóth, T., Lenková, M. (2015). Carrot (*Daucus carota* L. ssp. *sativus* (Hoffm.) Arcang.) as source of antioxidants. *Acta Agric. Slov.* 105, 303–311.
- Obia, O., & Emmanuel, F. D. (2025). Effect of Oral Administration of Common Pepper Types on the Liver Enzymes of Wistar Rats Fed with High-fat Diet. *East African Scholars Journal of Medical Sciences*. 8(3), 92-95.
- Ayeni, A., Abubakar, A., Aliyu, N., Uhomobhi, L., & Garba, I. (2019). 'Acute and sub-acute toxicity of the crude extracts of the aerial parts of *Daucus carota* L. in laboratory rats'. *Journal of Medicinal Plants for Economic Development*, 3(1), a69.
- Sambu, S., Hemaram, U., Murugan, R., & Alsofi, A. A. (2022). Toxicological and Teratogenic Effect of Various Food Additives: An Updated Review. *BioMed research international*, 2022, 6829409. <https://doi.org/10.1155/2022/6829409> (Retraction published Biomed Res Int. 2024 Jan 9;2024:9792751. doi: 10.1155/2024/9792751.)
- Warner, J. O. (2024). Artificial food additives: hazardous to long-term health?. *Archives of disease in childhood*, 109(11), 882–885. <https://doi.org/10.1136/archdischild-2023-326565>
- Yu, H., & Rhee, M. S. (2024). Potential of phytic acid in synergy with sodium chloride as a natural-borne preservative to inactivate *Escherichia coli* O157:H7 and inhibit natural microflora in fresh noodles at room temperature. *Current research in food science*, 9, 100868. <https://doi.org/10.1016/j.crfs.2024.100868>
- Baez, G., Chirio, M., Pisula, P., Seminario, E., Carasa, N., Philippi, R., Aroca-Martinez, G., & Musso, C. G. (2024). Hyponatremia and malnutrition: a comprehensive review. *Irish journal of medical science*, 193(2), 1043–1046. <https://doi.org/10.1007/s11845-023-03490-8>
- Puckett L. (2023). Renal and electrolyte complications in eating disorders: a comprehensive review. *Journal of eating disorders*, 11(1), 26. <https://doi.org/10.1186/s40337-023-00751-w>
- Obia, O., Odum, J. E., & Chuemere, A. N. (2018). Nephroprotective and antihyperlipidemic activity of honey in alloxan induced diabetic wistar rats. *International Journal of Biochemistry Research and Review*, 22(1), 1-7.
- Obia, O., & Eifuobhokhan, J. (2024). Effect of *Justicia carnea* leaf extract on plasma and fecal lipid profile in high-fat diet fed wistar rats.



International Journal of Health and Pharmaceutical Research, 9(4), 64-70.

22. Obia, O., Kalio, R. O., Tee, P. G. P., & Onyeso, G. (2025). Plasma Lipid Lowering Potential of Carrot (*Daucus carota*) Extract in Male Wistar Rats. *Asian Journal of Research in Medical and Pharmaceutical Sciences*, 14(1), 18-23.
23. Palmer, B. F., & Clegg, D. J. (2016). Achieving the Benefits of a High-Potassium, Paleolithic Diet, Without the Toxicity. *Mayo Clinic proceedings*, 91(4), 496–508.
24. Averin, E. E., Nikitin, A. E., Pozdnyak, A. O., Fedorova, E. L., Zhuk, V. S., Davydov, S. I., Fridman, I. L., Kompaniets, O. G., Kirpichnikova, N. V., Dudarenkova, M. R., Ginzburg, M. L., El Sharif, M. A., Martemyanova, E. G., & Sozykin, A. V. (2020). *Kardiologiya*, 60(2), 155–164. <https://doi.org/10.18087/cardio.2020.2.n972>
25. Goraya, N., Simoni, J., Jo, C., & Wesson, D. E. (2012). Dietary acid reduction with fruits and vegetables or bicarbonate attenuates kidney injury in patients with a moderately reduced glomerular filtration rate due to hypertensive nephropathy. *Kidney international*, 81(1), 86–93. <https://doi.org/10.1038/ki.2011.313>

